

NASA Contractor Report 191493

ICASE Report No. 93-37

191493
P-13

ICASE



ON CELL ENTROPY INEQUALITY FOR DISCONTINUOUS GALERKIN METHODS

Guangshan Jiang

Chi-Wang Shu

N94-11280

Unclass

G3/64 0181584

NASA Contract No. NAS1-19480
June 1993

Institute for Computer Applications in Science and Engineering
NASA Langley Research Center
Hampton, Virginia 23681-0001

Operated by the Universities Space Research Association



National Aeronautics and
Space Administration

Langley Research Center
Hampton, Virginia 23681-0001

(NASA-CR-191493) ON CELL ENTROPY
INEQUALITY FOR DISCONTINUOUS
GALERKIN METHODS Final Report
(ICASE) 13 P

ON CELL ENTROPY INEQUALITY FOR DISCONTINUOUS GALERKIN METHODS¹

Guangshan Jiang and Chi-Wang Shu

Division of Applied Mathematics

Brown University

Providence, Rhode Island 02912

ABSTRACT

We prove a cell entropy inequality for a class of high order discontinuous Galerkin finite element methods approximating conservation laws, which implies convergence for the one dimensional scalar convex case.

¹Research supported by ARO Grant DAAL03-91-G-0123, NSF Grant DMS-9211820, NASA Langley Grant NAG-1-1145 and Contract NAS1-19480 while the second author was in residence at ICASE, NASA Langley Research Center, Hampton, VA 23681-0001, and AFOSR Grant 93-0090.

1 Introduction

In [3] Cockburn and Shu defined a class of discontinuous Galerkin finite element methods for conservation laws (the multi-dimensional case was discussed in [4]):

$$u_t + \operatorname{div} f(u) = 0 \quad (1.1)$$

The entropy solution of (1.1) also satisfies

$$U(u)_t + \operatorname{div} F(u) \leq 0 \quad (1.2)$$

in the distribution sense, for any convex function $U(u)$ and consistent entropy flux $F(u)$ satisfying $F'(u) = U'(u)f'(u)$. The scheme is obtained in the following way: first a triangulation \mathcal{T}_h is chosen. In one space dimension \mathcal{T}_h is just a collection of subintervals $I_j = (x_{j-\frac{1}{2}}, x_{j+\frac{1}{2}})$, which are not necessarily of the same length. We will use the notation $\Delta x_j = x_{j+\frac{1}{2}} - x_{j-\frac{1}{2}}$, $h = \max_j \Delta x_j$ and $x_j = \frac{1}{2}(x_{j-\frac{1}{2}} + x_{j+\frac{1}{2}})$. In two and more space dimensions \mathcal{T}_h is a collection of triangles, tetrahedrons or other simple geometric objects. The solution space V_h is defined as the collection of all piecewise polynomials of degree up to r for a $(r+1)$ -th order method. The functions in V_h are allowed to have discontinuities across element interface. The conservation law (1.1) is multiplied by a test function $v \in V_h$, integrated over an element $K \in \mathcal{T}_h$, and formally integrated by parts to shift the spatial derivatives from $f(u)$ to v . The result is:

$$\int_K u_t(x, t) v(x) dx + \sum_{e \in \partial K} \int_e f(u(x, t)) \cdot n v(x) ds - \int_K f(u(x, t)) \cdot \operatorname{grad} v(x) dx = 0. \quad (1.3)$$

where ∂K is the boundary of the element K and n is the unit outward normal vector. This procedure up to now is just the standard discontinuous Galerkin method. Two features from the finite difference methodology are then used in [3], [4]: one is the monotone or E flux (approximate Riemann solver for systems) to define the trace of the physical flux $f(u)$ at the cell interface; the other is a local nonlinear limiter which limits the values of u at the

cell interface by the differences of the means of u over elements. With these two ingredients, it was proven in [3] and [4] that the schemes are formally high order accurate, measured by local truncation errors in smooth regions including at extrema, are total variation bounded for one space dimension, and are maximum norm bounded for any space dimensions. The only thing missing is the entropy condition: in [3], we were able to prove entropy consistency for the square entropy $U(u) = \frac{u^2}{2}$ for one dimensional convex $f(u)$ with a h -independent modification to the scheme, following the idea of Osher [12]. We were also able to prove entropy consistency for all convex entropies with a h -dependent modification to the scheme. The h -dependent limiters make the proof of high order schemes easy, but it is not very desirable for practical computations, because it usually limits the slope near the discontinuities stronger than necessary, and essentially flattens the solution there to piecewise constants for fine mesh. It also destroys the self-similarity of the scheme.

Entropy condition seems difficult to prove for high order finite difference schemes. Osher and Tadmor proved [13] that finite difference schemes (which evolve only the mean) which satisfy cell entropy inequalities for all convex entropies can be at most first order accurate. Even for one entropy inequality (say for the square entropy), for one space dimension and for convex $f(u)$, the proof is extremely elusive if one does not modify the scheme. Osher [11] and Nessyahu and Tadmor [10] were able to get such cell entropy inequalities, for the square entropy and for the second order MUSCL scheme, with some h -independent modifications; Yang [14] was able to prove convergence of unmodulated second order MUSCL scheme using a global analysis rather than relying solely on cell entropy inequalities; Lions and Souganidis [8] proved convergence of second order MUSCL scheme for steady state Hamilton-Jacobi equations and conservation laws. There are also many results which prove entropy consistency and/or convergence using h -dependent limiters or modifications for high order schemes. For example Coquel and LeFloch [7] for finite difference; Johnson, Szepessy and Hansbo [9] for streamline diffusion finite element; Cockburn, Coquel and LeFloch [5] and Cockburn and Gremaud [6] for high order finite volume, streamline diffusion or discontinuous

Galerkin schemes with h -dependent “shock capturing” terms. These results are usually more general (multi space dimensions, non-convex fluxes, etc.). However, as we have indicated before, h -dependent limiters or modifications should be avoided for practical computations if possible. More recently, Bouchut and Perthame [2] obtained a second order one dimensional scheme which is consistent with all entropy conditions and does not use h -dependent limiter. A key ingredient of [2] is to evolve both the mean and the slope and use the whole function (not just the mean) to obtain cell entropy inequalities. Discontinuous Galerkin methods also fall into this category (evolving the whole polynomial in a cell rather than just the mean), and we use this fact strongly in this paper.

In Section 2 we prove a cell entropy inequality using the square entropy $U(u) = \frac{u^2}{2}$ for the unmodulated semi-discrete discontinuous Galerkin method of [3], [4]. The proof is remarkably simple and does not even use any nonlinear limiters. It works thus for any spatial order of accuracy. As far as we know, all the previous cell entropy inequalities without h -dependent limiters must restrict the slope of a function by $\frac{1}{\Delta x_j} \minmod(\Delta_+ u_j, \Delta_- u_j)$ (which means that the difference at the interface, $u_{j+\frac{1}{2}}^+ - u_{j+\frac{1}{2}}^-$, must be of the same sign as that of $u_{j+1} - u_j$, i.e., no “sawtooth” is allowed in the reconstruction), hence cannot be higher than second order accurate. The result in this paper illustrates the potential of discontinuous Galerkin methods, or equivalent Hermite type finite difference/finite volume type methods, which evolve the whole polynomial in the cell rather than just the mean. Time discretization is discussed in Section 3.

2 Cell Entropy Inequality for the Square Entropy

The discontinuous Galerkin scheme in one space dimension, defined in [3], in its semi-discrete form without slope limiting, is the following: Find $u(\cdot, t) \in V_h$, such that, for all $v \in V_h$ and all subinterval I_j :

$$\int_{I_j} u_t(x, t) v(x) dx + h_{j+\frac{1}{2}}(t) v(x_{j+\frac{1}{2}}^-) - h_{j-\frac{1}{2}}(t) v(x_{j-\frac{1}{2}}^+) - \int_{I_j} f(u(x, t)) v_x(x) dx = 0 \quad (2.1)$$

Here, $h_{j+\frac{1}{2}}(t) = h\left(u(x_{j+\frac{1}{2}}^-, t), u(x_{j+\frac{1}{2}}^+, t)\right)$ is a Lipschitz continuous monotone flux (i.e. h is non-decreasing in the first argument and non-increasing in the second argument), or more generally an E -flux as defined by Osher [11]:

$$\left(h(u^-, u^+) - f(u)\right)(u^+ - u^-) \leq 0 \quad (2.2)$$

for all u between u^- and u^+ . Some examples of the commonly used monotone fluxes can be found in, e.g., [3].

If we take $v(x) = u(x, t)$ in (2.1), we get

$$\int_{I_j} \left(\frac{u^2(x, t)}{2} \right)_t dx + h_{j+\frac{1}{2}}(t)u(x_{j+\frac{1}{2}}^-, t) - h_{j-\frac{1}{2}}(t)u(x_{j-\frac{1}{2}}^+, t) - \int_{I_j} f(u(x, t))u_x(x, t)dx = 0 \quad (2.3)$$

We define

$$g(u) = \int^u f(u)du \quad (2.4)$$

and rewrite (2.3) as

$$\int_{I_j} \left(\frac{u^2(x, t)}{2} \right)_t dx + \hat{F}_{j+\frac{1}{2}}(t) - \hat{F}_{j-\frac{1}{2}}(t) + A_j(t) = 0 \quad (2.5)$$

where

$$\hat{F}_{j+\frac{1}{2}}(t) = h_{j+\frac{1}{2}}(t)u(x_{j+\frac{1}{2}}^-, t) - g(u(x_{j+\frac{1}{2}}^-, t)) \quad (2.6)$$

is consistent with the entropy flux for the square entropy:

$$F(u) = \int^u f'(u)udu = f(u)u - \int^u f(u)du = f(u)u - g(u), \quad (2.7)$$

and

$$A_j(t) = -h_{j-\frac{1}{2}}(t) \left(u(x_{j-\frac{1}{2}}^+, t) - u(x_{j-\frac{1}{2}}^-, t) \right) + g(u(x_{j-\frac{1}{2}}^+, t)) - g(u(x_{j-\frac{1}{2}}^-, t))$$

$$= -\left(h_{j-\frac{1}{2}}(t) - f(\xi)\right) \left(u(x_{j-\frac{1}{2}}^+, t) - u(x_{j-\frac{1}{2}}^-, t)\right) \geq 0 \quad (2.8)$$

where we have used the mean value theorem and the definition (2.4) of $g(u)$ in the second equality, ξ is between $u(x_{j-\frac{1}{2}}^-, t)$ and $u(x_{j-\frac{1}{2}}^+, t)$, and the last inequality is due to the property of E-flux (2.2).

We have thus proven the cell entropy inequality

$$\int_{I_j} \left(\frac{u^2(x, t)}{2} \right)_t dx + \hat{F}_{j+\frac{1}{2}}(t) - \hat{F}_{j-\frac{1}{2}}(t) \leq 0 \quad (2.9)$$

for the square entropy $U(u) = \frac{u^2}{2}$. Notice that we do not need any nonlinear limiting at this stage. However, nonlinear limiting as introduced in [3] and [4] will not destroy this cell entropy inequality (see next section). The cell entropy inequality (2.9) trivially implies L^2 stability of the scheme (again without even using the nonlinear limiting):

$$\frac{d}{dt} \int_{-\infty}^{\infty} \left(\frac{u^2(x, t)}{2} \right) dx \leq 0 \quad (2.10)$$

but it is much stronger. For example, if $f(u)$ is convex and we use the nonlinear limiting [3] to obtain a total variation boundedness for the solution, we will have convergence towards the unique entropy solution.

The same entropy inequality can be obtained for multi-space dimensions with arbitrary triangulations:

$$\int_K \left(\frac{u^2(x, t)}{2} \right)_t dx + \sum_{e \in \partial K} \int_e F_{e,K}(x, t) ds \leq 0 \quad (2.11)$$

where $F_{e,K}$ is consistent with $F \cdot n_K$ for the entropy flux F in (1.2) and the outward normal n_K , and $F_{e,K} = -F_{e,K'}$ for the two neighboring elements $K \cap K' = \{e\}$ (conservation). We omit the derivation detail since it parallels that for the one dimensional case.

We have the following two remarks:

(1). If we try to do the same estimate for a general convex entropy $U(u)$, we can obtain exactly the same cell entropy inequality module an interpolation error term:

$$- \int_{I_j} (u(x, t)_t + f(u(x, t))_x) (U'(u(x, t)) - \Pi U'(u(x, t))) dx \quad (2.12)$$

where Πv is a projection into the space V_h which interpolates at the two end points of I_j . This motivates the following h -dependent modification to the scheme, which is similar to the “shock capturing term” added to the streamline diffusion method in Johnson, Szepessy and Hansbo [9] and in Cockburn and Gremaud [6]:

$$\begin{aligned} \int_{I_j} u_t(x, t) v(x) dx &+ h_{j+\frac{1}{2}}(t) v(x_{j+\frac{1}{2}}^-) - h_{j-\frac{1}{2}}(t) v(x_{j-\frac{1}{2}}^+) - \int_{I_j} f(u(x, t)) v_x(x) dx \\ &+ Ch \int_{I_j} |u_t(x, t) + f(u(x, t))_x| \frac{u_x(x, t)}{|u_x(x, t)|} v_x(x) dx = 0 \end{aligned} \quad (2.13)$$

where C is a suitable positive constant. This, together with a L_∞ bound which can be obtained by using nonlinear limiters [3], will give us a cell entropy inequality for arbitrary convex entropy at least for the $r = 1$ (second order) case, hence convergence for any nonconvex flux $f(u)$ in this case. For general r , more h -dependent modification is needed. Notice that by adding this modification, the *formal* order of accuracy of the scheme is not changed: $|u_t(x, t) + f(u(x, t))_x|$ is just the local truncation error. A similar argument as in [9] shows that the modification does not destroy convergence towards weak solutions (conservation).

(2). The so-called discontinuous Galerkin method can also be recasted as a finite difference scheme (Hermite type: where one evolves both the mean and the slope, maybe more). For example, the second order case is just the following scheme for the mean $\bar{u}_j(t)$ and the slope $s_j(t)$ where $u(x, t) = \bar{u}_j(t) + s_j(t)(x - x_j)$ in cell I_j (see [3]):

$$\begin{aligned} \frac{d}{dt} \bar{u}_j(t) &= -\frac{1}{\Delta x_j} [\hat{f}_{j+\frac{1}{2}}(t) - \hat{f}_{j-\frac{1}{2}}(t)] \\ \frac{d}{dt} s_j(t) &= -\frac{6}{\Delta x_j^2} [\hat{f}_{j+\frac{1}{2}}(t) + \hat{f}_{j-\frac{1}{2}}(t)] + \frac{12}{\Delta x_j^3} \int_{I_j} f(\bar{u}_j(t) + s_j(t)(x - x_j)) dx \end{aligned} \quad (2.14)$$

where

$$\hat{f}_{j+\frac{1}{2}}(t) = h \left(\bar{u}_j(t) + \frac{\Delta x_j}{2} s_j(t), \bar{u}_{j+1}(t) - \frac{\Delta x_{j+1}}{2} s_{j+1}(t) \right) \quad (2.15)$$

We have thus in effect proved cell entropy inequalities for such high order Hermite type finite difference schemes without using the help of any nonlinear limiting. Of course, to get convergence one must use the nonlinear limiting to obtain L_∞ and/or total variation bounds.

3 Time Discretization

We discretize (2.1) in time by the following class of methods:

$$\begin{aligned} \int_{I_j} \frac{u^{n+1}(x) - u^n(x)}{\Delta t} v(x) dx &+ h_{j+\frac{1}{2}}^{n+\theta} v(x_{j+\frac{1}{2}}^-) - h_{j-\frac{1}{2}}^{n+\theta} v(x_{j-\frac{1}{2}}^+) \\ &- \int_{I_j} f(u^{n+\theta}(x)) v_x(x) dx = 0 \end{aligned} \quad (3.1)$$

where

$$\begin{aligned} u^{n+\theta}(x) &= (1 - \theta)u^n(x) + \theta u^{n+1}(x) \\ h_{j+\frac{1}{2}}^{n+\theta} &= h \left(u^{n+\theta}(x_{j+\frac{1}{2}}^-), u^{n+\theta}(x_{j+\frac{1}{2}}^+) \right) \end{aligned} \quad (3.2)$$

For $\theta = 0$ this is the Euler forward discretization; for $\theta = 1$ this is Euler backward; and for $\theta = \frac{1}{2}$ this is Crank-Nicolson.

If we take $v(x) = u^{n+\theta}(x)$, we obtain just as before

$$\int_{I_j} \frac{u^{n+1}(x) - u^n(x)}{\Delta t} u^{n+\theta}(x) dx + \hat{F}_{j+\frac{1}{2}}^{n+\theta} - \hat{F}_{j-\frac{1}{2}}^{n+\theta} \leq 0 \quad (3.3)$$

where

$$\hat{F}_{j+\frac{1}{2}}^{n+\theta} = h_{j+\frac{1}{2}}^{n+\theta} u^{n+\theta}(x_{j+\frac{1}{2}}^-) - g \left(u^{n+\theta}(x_{j+\frac{1}{2}}^-) \right) \quad (3.4)$$

with $g(u)$ defined by (2.4). We can rewrite (3.3) as

$$\int_{I_j} \frac{(u^{n+1}(x))^2 - (u^n(x))^2}{2\Delta t} dx + \hat{F}_{j+\frac{1}{2}}^{n+\theta} - \hat{F}_{j-\frac{1}{2}}^{n+\theta} + \left(\theta - \frac{1}{2}\right) \int_{I_j} \frac{(u^{n+1}(x) - u^n(x))^2}{\Delta t} dx \leq 0 \quad (3.5)$$

Thus a sufficient condition to get the cell entropy inequality

$$\int_{I_j} \frac{(u^{n+1}(x))^2 - (u^n(x))^2}{2\Delta t} dx + \hat{F}_{j+\frac{1}{2}}^{n+\theta} - \hat{F}_{j-\frac{1}{2}}^{n+\theta} \leq 0 \quad (3.6)$$

is just $\theta \geq \frac{1}{2}$, i.e. implicit schemes from Crank-Nicolson to Backward Euler.

Up to now we have not considered the nonlinear *minmod* type limiters in [3] and [4]. These limiters will render the scheme total variation stable (in one space dimension) or L_∞ stable (in multi space dimensions). We have to make sure that these limiters do not destroy the cell entropy inequality (3.6). This turns out to be a simple issue: suppose $\tilde{u}^{n+1}(x)$ is the solution obtained from the scheme (3.1) without limiter, then projected to get the solution at time level $n+1$ by some *minmod* type limiter $u^{n+1}(x) = P\tilde{u}^{n+1}(x)$ (this is the procedure adopted in [3] and [4]). The above derivation for the unlimited scheme will give (3.6) for \tilde{u}^{n+1} :

$$\int_{I_j} \frac{(\tilde{u}^{n+1}(x))^2 - (u^n(x))^2}{2\Delta t} dx + \hat{F}_{j+\frac{1}{2}}^{n+\theta} - \hat{F}_{j-\frac{1}{2}}^{n+\theta} \leq 0 \quad (3.7)$$

hence a sufficient condition to get the cell entropy inequality (3.6) for the limited (projected) solution u^{n+1} is to require the projection P to satisfy

$$\int_{I_j} (Pw(x))^2 dx \leq \int_{I_j} (w(x))^2 dx \quad (3.8)$$

for all polynomial $w(x)$ of degree up to r (recall that $r+1$ is the order of the scheme), i.e. P does not increase the L_2 norm of any r -th order polynomial in cell I_j . Notice that this is the idea used in [2]. An easy way to ensure condition (3.8) is to write $w(x)$ and $Pw(x)$ as expansions of (scaled) Legendre polynomials $p_l(x)$ as in [3]: $w(x) = \sum_{l=0}^r a_l p_l(x)$ and $Pw(x) = \sum_{l=0}^r b_l p_l(x)$. A simple sufficient condition to ensure (3.8) is now just $|b_l| \leq |a_l|$ for all l , which is easily checked to be correct for the P^1 case ($r=1$) in [3] and also for the general P^r case with similar *minmod* limiters on a_l (see also [1] for such limiters).

It is also possible to get cell entropy inequalities for certain explicit time discretizations. For example, with leap-frog time discretization:

- [5] B. Cockburn, F. Coquel and F. LeFloch, *An error estimate for finite volume methods for multidimensional conservation laws*, Math. Comput., to appear.
- [6] B. Cockburn and P. Gremaud, *An error estimate for finite element methods for scalar conservation laws*, SIAM J. Numer. Anal., submitted.
- [7] F. Coquel and F. LeFloch, *Convergence of finite difference schemes for conservation laws in several space dimensions: the corrected antidiffusive flux approach*, Math. Comput., v57 (1991), pp.169-210.
- [8] P. Lions and P. Souganidis, *Convergence of MUSCL type methods for scalar conservation laws*, C. R. Acad. Sci. Paris, v311 (1990), pp.259-264.
- [9] C. Johnson, A. Szepessy and P. Hansbo, *On the convergence of shockcapturing streamline diffusion finite element methods for hyperbolic conservation laws*, Math. Comput., v54 (1990), pp.107-129.
- [10] H. Nessyahu and E. Tadmor, *Non-oscillatory central differencing for hyperbolic conservation laws*, J. Comput. Phys., v87 (1990), pp.408-463.
- [11] S. Osher, *Riemann solvers, the entropy condition, and difference approximations*, SIAM J. Numer. Anal., v21 (1984), pp.217-235.
- [12] S. Osher, *Convergence of generalized MUSCL schemes*, SIAM J. Numer. Anal., v22 (1985), pp.947-961.
- [13] S. Osher and E. Tadmor, *On the convergence of difference approximations to scalar conservation laws*, Math. Comput., v50 (1988), pp.19-51.
- [14] H. Yang, *Extremum tracking and convergence of semi-discrete TVD schemes*, IMA Preprint Series # 697, September 1990, University of Minnesota.

$$\int_{I_j} \frac{u^{n+1}(x) - u^{n-1}(x)}{2\Delta t} v(x) dx + h_{j+\frac{1}{2}}(t)^n v(x_{j+\frac{1}{2}}^-) - h_{j-\frac{1}{2}}(t)^n v(x_{j-\frac{1}{2}}^+) - \int_{I_j} f(u^n(x)) v_x(x) dx = 0 \quad (3.9)$$

we can obtain, by taking $v(x) = u^n(x)$:

$$\int_{I_j} \frac{u^{n+1}(x)u^n(x) - u^n(x)u^{n-1}(x)}{2\Delta t} dx + \hat{F}_{j+\frac{1}{2}}^n - \hat{F}_{j-\frac{1}{2}}^n \leq 0. \quad (3.10)$$

4 Concluding Remarks

We have shown that the discontinuous Galerkin method, or equivalent Hermite type finite difference method, which evolves the whole polynomial inside a cell, satisfies a cell entropy inequality for the square entropy in multi space dimensions. Limiters to enforce total variation or maximum norm stability can be designed not to destroy this cell entropy inequality. Time discretization is also discussed.

References

- [1] R. Biswas, K. Devine and J. Flaherty, *Parallel, adaptive finite element methods for conservation laws*, Appl. Numer. Math., to appear.
- [2] F. Bouchut, C. Bourdarias and B. Perthame, *Un exemple de methode MUSCL satisfaisant toutes les inegalites d'entropie numeriques*, preprint, to be submitted to C.R. Acad. Sci. Paris.
- [3] B. Cockburn and C.-W. Shu, *TVB Runge-Kutta local projection discontinuous Galerkin finite element method for conservation laws II: general framework*, Math. Comput., v52 (1989), pp.411-435.
- [4] B. Cockburn, S. Hou and C.-W. Shu, *The Runge-Kutta local projection discontinuous Galerkin finite element method for conservation laws IV: the multidimensional case*, Math. Comput., v54 (1990), pp.545-581.

REPORT DOCUMENTATION PAGE			Form Approved OMB No 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE June 1993	3. REPORT TYPE AND DATES COVERED Contractor Report		
4. TITLE AND SUBTITLE ON CELL ENTROPY INEQUALITY FOR DISCONTINUOUS GALERKIN METHODS		5. FUNDING NUMBERS C NAS1-19480 WU 505-90-52-01		
6. AUTHOR(S) Guangshan Jiang Chi-Wang Shu				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Institute for Computer Applications in Science and Engineering Mail Stop 132C, NASA Langley Research Center Hampton, VA 23681-0001		8. PERFORMING ORGANIZATION REPORT NUMBER ICASE Report No. 93-37		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Langley Research Center Hampton, VA 23681-0001		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA CR-191493 ICASE Report No. 93-37		
11. SUPPLEMENTARY NOTES Langley Technical Monitor: Michael F. Card Final Report To appear in Mathematics of Computation				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Unclassified - Unlimited Subject Category 64			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) We prove a cell entropy inequality for a class of high order discontinuous Galerkin finite element methods approximating conservation laws, which implies convergence for the one dimensional scalar convex case.				
14. SUBJECT TERMS conservation law, discontinuous Galerkin, entropy condition, convergence			15. NUMBER OF PAGES 12	
			16. PRICE CODE A03	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	